

SOUTHWEST FISHERIES CENTER

NATIONAL MARINE FISHERIES SERVICE

SOUTHWEST FISHERIES CENTER

P.O. BOX 271

LA JOLLA, CA 92038

LIBRARY

SEP 6 1988

National Marine Fisheries Service
2570 Dole Street
Honolulu, Hawaii

JULY 1988

PLAN FOR COLLECTION OF FISHERIES DATA FROM U.S. TUNA PURSE SEINERS FISHING IN THE SOUTH PACIFIC

by

Atilio L. Coan Jr., Norman Bartoo and Gary Sakagawa

This Administrative Report is issued as an informal document to ensure prompt dissemination of preliminary results, interim reports and special studies. We recommend that it not be abstracted or cited.

PLAN FOR COLLECTION OF FISHERIES DATA FROM U.S. TUNA
PURSE SEINERS FISHING IN THE SOUTH PACIFIC

Atilio L. Coan Jr.

Norman Bartoo

and

Gary Sakagawa

National Marine Fisheries Service
Southwest Fisheries Center
La Jolla, California

July 1988

CONTENTS

	Page
INTRODUCTION	1
PURPOSE OF SAMPLING	1
LOGBOOK DATA	2
UNLOADING WEIGHTS	2
SIZE-FREQUENCY AND SPECIES COMPOSITION DATA	3
Selection of Wells	4
Sampling Location	4
Size-sorted Unloading	4
Size and Species Composition Sampling	4
BIOLOGICAL SAMPLES	5
TAG RECAPTURE INFORMATION	5
VESSEL INFORMATION	6
SAMPLING PORTS	6
DISTRIBUTION OF DATA	6
ACKNOWLEDGMENTS	6
LITERATURE CITED	7

LIST OF FIGURES

	Page
Figure 1. South Pacific Treaty region and statistical areas used for sampling of catches and fish sizes	8
Figure 2. Purse Seine Vessel Catch Report (PSVCR) Form used to record daily fishing information. The form is completed by the vessel captains and is submitted to the FFA as required under the Treaty	9
Figure 3. Logbook Abstracting (LA) Form used to record information for selecting wells for sampling of fish sizes	10
Figure 4. Purse Seine Unloading Logsheet (PSUL) used to record cannery unloading weights. The form is completed by the vessel captain and is submitted to the FFA as required under the Treaty	11
Figure 5. Trip/Unloading (TU) Form used to compare vessel unloading weights and hailed (estimated at time of catch) weights.	12
Figure 6. Length-Frequency Sampling Status Board used to monitor the size sampling by month-area strata.	13
Figure 7. Vessel Length-Frequency Sampling Plan used to prioritize size sampling of individual vessel's catches.	14
Figure 8. Length-Frequency (LF) Form used to record sizes of fish, catches and catch location by vessel well.	15-16
Figure 9. Species Composition (SC) Form used to record proportions of each species in the catch by vessel well	17
Figure 10. Tag Recovery Record used to record weight, length, catch area and other information for tag returns	18
Figure 11. Monthly summary of tags and reward payments form used to account for distribution of rewards.	19

PLAN FOR COLLECTION OF FISHERIES DATA FROM U.S. TUNA PURSE SEINERS FISHING IN THE SOUTH PACIFIC

Atilio L. Coan Jr., Norman Bartoo and Gary Sakagawa

INTRODUCTION

A South Pacific Regional Tuna Treaty that allows licensing of U.S. tuna purse seiners to fish in the 200-mile economic zones of sixteen South Pacific island states¹ was ratified in June 1988 by the U.S. Congress. In preparation for the ratification, the Southwest Fisheries Center (SWFC) of the National Marine Fisheries Service (NMFS) developed plans to implement provisions of the Treaty with respect to port sampling of the catches made by licensed purse seiners.

A meeting was held on January 6, 1988 at the SWFC in La Jolla, California, to coordinate the planning among the major parties, NMFS Southwest Region (SWR) and the Forum Fisheries Agency (FFA), an agency designated by the South Pacific Island Governments to handle administration of the Treaty. Attending the meeting were representatives of the SWFC, SWR, FFA, and South Pacific Commission (SPC). The SPC representative participated as a technical consultant to the FFA delegation. The participants agreed that NMFS would conduct the port sampling for all parties and follow certain sampling procedures for the 1988 season. The rationale and sampling procedures that will be used are described in this document.

PURPOSE OF SAMPLING

Between 30 and 40 U.S. purse seiners are expected to be licensed to fish for yellowfin (*Thunnus albacares*) and skipjack (*Katsuwonus pelamis*) tunas in the South Pacific Treaty region (Figure 1) during 1988. In compliance with requirements of the Treaty, NMFS representatives will collect five types of fisheries statistics from these vessels and their landed catches: (1) logbook data on fishing operations, (2) unloading weights by species, (3) size frequency of landed catch, (4) species composition of landed catch, and (5) biological material.

¹South Pacific states adhering to the Treaty are Australia, Cook Island, Federated States of Micronesia, Fiji, Kiribati, Marshall Island, Nauru, New Zealand, Niue, Palau, Papua New Guinea, Solomon Island, Tonga, Tuvalu, Vanuatu, Western Samoa.

Additionally, representatives will collect secondary information on vessel operations and methods and tagged fish that are returned.

LOGBOOK DATA

Purse Seine Vessel Catch Report Forms (PSVCR Form, Figure 2) will be distributed, by NMFS representatives, to licensed U.S. purse seiners at the beginning of the fishing season. Vessel captains are required to complete this form by making daily entries when fishing in the Treaty area. The completed form will be collected by a NMFS representative upon a vessel's arrival in port for unloading or transshipment (e.g. end of fishing trip).

Three separate logs are usually maintained by U.S. vessels: captain's fishing log, navigator's log and engineer's log. The representative will use these logs and the PSVCR Form to select vessel wells that will be sampled for size and species composition. The engineer's log provides locations of catches from individual sets in wells (e.g. small and large fish from the same set that are separated and stored in different wells) and may be useful in identifying the location of a few tons of incidental species. The navigator's and captain's logs are useful for obtaining locations of single sets, vessel positions, and daily vessel activity.

To facilitate selection of wells to be sampled, pertinent information from the three vessel logs will be abstracted to a separate Logbook Abstracting Form (LA Form, Figure 3). Abstracting is a necessary precursor to selecting wells for sampling of the catch as well as in consolidating questions for later review with the vessel's captain and crew.

UNLOADING WEIGHTS

Large metal containers are used to unload tuna at ports with cannery facilities. The containers are filled on the vessel's deck, hoisted to the dock, weighed and stored in cannery holding facilities. The unloading weights by species are recorded by cannery personnel and should be routinely collected by the NMFS representative.

The Purse Seine Unloading Logsheet (PSUL Form, Figure 4) will be collected from the vessel captain, who is required to complete one for each unloading. The NMFS representative should inspect this form for any at-sea transfers of fish, as they may affect selection of wells for size sampling. At-sea transfers are usually noted on the PSVCR Form or in the captain's log.

Catches on the PSVCR Form are estimates or "hailed weights." They are not the same as the unloading weights from the canneries. After a vessel has unloaded, the NMFS representative will enter total catch by species from the PSVCR Form and unloading weights from the PSUL Form on the Trip/Unloading Form (TU Form, Figure 5) and compare the two weights. Differences between these weights can be due to catches not entered for some sets, erroneous species identification, unrecorded catches discarded or transferred at sea, or catches unloaded at an earlier date. If the difference between logged total catch and unloading weight is greater than 25%, the vessel captain should be consulted to identify the source of discrepancy.

SIZE-FREQUENCY AND SPECIES COMPOSITION DATA

The NMFS representative will schedule sampling of vessel wells in order to maximize the likelihood of obtaining samples that meet requirements of the sampling plan. The schedule will be dictated primarily by the cannery's unloading schedule, and should incorporate LA Form information on time and area of catch, and storage of sets in wells. Using this information, the NMFS representative will prepare and maintain a Length-Frequency Sampling Status Board (Figure 6), to keep track of samples taken to date and prepare a Vessel Length-Frequency Sampling Plan (Figure 7) that identifies samples to be taken from each vessel.

As vessels are unloaded, NMFS representatives will take length-frequency and species composition samples from the unloaded catch. The objective of length-frequency sampling is to compile size information for yellowfin, bigeye and skipjack tunas from various time-area strata of the catch. The objective of species composition sampling is to accurately estimate the proportions of yellowfin, skipjack and bigeye tunas in the reported catches.

Sampling will be stratified by month and by area. Five area strata have been established (Figure 1) from inspection of past catch records for uniformity of fish sizes and clustering of fishing effort. The area strata maximize the chances for sampling fish from unique month-area catches.

A minimum of 13 yellowfin, 13 bigeye, and 10 skipjack length-frequency samples per month-area strata will be collected. Samples of 100 fish will be collected for species composition and of 50 fish for length frequencies (Hennemuth 1957). Samples will be drawn from as many different vessels and wells as possible to maximize the accuracy of the sampling (Coan and Bartoo 1984). If time permits, more samples from strata already meeting the minimum sample size criterion will be scheduled (Wetherall and Yong, In Press).

Species composition samples will be taken whenever a mixture of fish species is encountered. Usually, skipjack are separated

from yellowfin at time of unloading. Consequently, only landings labeled yellowfin or bigeye normally need sampling to determine the amount of each species mixed in the landing.

Selection of Wells

The NMFS representative will sample wells that contain catches from one month-area stratum before sampling wells that contain catches from multiple strata. If only wells with catches from multiple strata are available, wells with catches that are layered by strata will be selected (e.g., bottom of the well, top of well, etc.). Sampling a particular layer will require waiting until the desired portion of the well is unloaded before drawing the sample.

Sampling Location

The best location for drawing samples is aboard the vessel close to the well being unloaded. As fish are unloaded, a random sample of each species can be drawn.

If sampling aboard the vessel is not permitted, then sampling should be done on the dock during offloading of the containers and before fish are stored or processed. Random samples are drawn from the containers. Information on species, well and size group (if sorting by size occurs -- see below) should be collected. Sampling should be done before the fish are size sorted whenever possible.

Size-sorted Unloading

At times the unloading crews will sort fish by size groups (small, medium, large). This size sorting makes sampling more complex. Fifty fish samples for each species from each size group must be drawn. If mixed species are encountered, species composition samples must also be taken for each size group. The unloading weight for each size group by well and for the entire vessel must be noted on the sampling form. The unloading weight by size group for the well can be estimated by counting the number of buckets unloaded for each size group. Unloading weights by size group for the entire vessel's load can be obtained from the canneries.

Size and Species Composition Sampling

Wells selected for sampling should be examined for mixed species. If mixing is detected, 100 fish will be randomly drawn and the size and species of each fish recorded. Sizes of fish must be recorded in order to determine the proportion of each species by weight in the sampled well. A Length-Frequency Form (LF Form, Figure 8) should be used to record the sizes of fish

and a Species Composition Form (SC Form, Figure 9) to record the numbers of each species. After the 100 fish are drawn and if sufficient numbers of fish are present, extra fish are drawn to complete a 50 fish length-frequency sample for yellowfin, bigeye or skipjack. Fifty fish samples of other species in the well can be drawn if sufficient quantities of these fish exist.

If mixing of species in the well is not detected, then species composition samples will not be taken. Instead, 50 fish will be randomly drawn from the selected well, measured for length and recorded on the LF Form.

Fish will be measured for fork length (tip of snout to fork in tail) to the nearest centimeter whenever possible. If fish are too large to handle, predorsal lengths (snout to anterior base of the dorsal fin) can be taken; fork-length measurements, however, are preferred. The type of length measurement must be noted on the LF Form.

BIOLOGICAL SAMPLES

Depending on time constraints, the NMFS representatives will collect biological samples from fish measured for length frequency or in the cannery at the butchering line. Exact sampling procedures will depend on the type of material (otoliths, livers, etc.) requested by researchers.

Biological samples require some form of preservation before shipment to researchers. At times, preservation requires the use of chemicals that the canneries will not allow on the premises. Also, whole fish samples usually cannot be stored safely at the cannery. Availability of laboratory and storage facilities away from cannery operations will determine the number and types of biological samples that can be collected.

TAG RECAPTURE INFORMATION

Tags are generally found as fish are removed from the wells or as fish are butchered and processed on the canning line. The NMFS representative will be responsible for collecting tag return information. Upon receipt of a tag the finder is paid a reward and all relevant information such as the name of the catcher vessel, date landed, etc. is recorded on the Tag Recovery Record (Figure 10). A summary of all tags received and rewards made will be maintained (Figure 11).

VESSEL INFORMATION

Vessel and fishing operations data may occasionally be required. NMFS representatives will be available to collect such data as time permits.

SAMPLING PORTS

The number of NMFS representatives at each landing port will depend on the volume and frequency of landings. During 1986, U.S. purse seiners fishing in the western tropical Pacific made 93 separate landings in Pago Pago, American Samoa, 35 in Guam/Tinian, 5 in Thailand, 3 in the Philippines, 3 in Indonesia and 1 in Australia.

For the 93 landings in Pago Pago, only seven times during the year did 2 vessels land on the same day; most of the time only one vessel landed per day. Under optimum conditions, the minimum time required to abstract one logbook is 1 hour and to obtain one size sample is two hours (includes species composition). Taking these and other considerations such as vacations and sick leave into account, a minimum of two representatives will be required for Pago Pago. Sampling in Guam/Tinian will probably require one or two representatives; however, since the number of landings in Guam/Tinian varies depending on transshipment costs to the vessels, coverage needs to be evaluated yearly. Landings at the other ports appear to be too infrequent to warrant on-site sampling by permanent staff; however, other arrangements for sampling could be made.

DISTRIBUTION OF DATA

Port sampling data will be prepared by the NMFS representatives: one copy retained by the field office, one sent to the SWFC and one copy (the original PSVCR Form and PSUL Form) sent to the FFA. The SWFC will enter the data to computer databases and run edit programs to screen the data for inconsistent information. After editing, the data on computer tapes will be made available to the FFA.

ACKNOWLEDGMENTS

We would like to acknowledge the assistance of Mike Hinton, Inter-American Tropical Tuna Commission, for providing general information on sampling of Pacific tunas and Frank Alverson, LMR Fisheries Research, Inc., for information on fleet movements and fish sizes in catches from the South Pacific.

LITERATURE CITED

- Coan, A. L. and N. Bartoo. 1984. Effects of sample size on the accuracy of length-frequency sampling of tunas transshipped to Puerto Rico. ICCAT Coll. Vol. of Sci. Paps. (SCRS-1983), Vol. XX, Madrid, Spain. p. 68-74.
- Hennemuth R.C. 1957. An analysis of methods of sampling to determine the size composition of commercial landings of yellowfin tuna (*Neothunnus macropterus*) and skipjack tuna (*Katsuwonus pelamis*). [In Engl. and Span.] Inter-Am. Trop. Tuna Comm., Bull. 2:174-243.
- Wetherall J. A. and M. Y. Y. Yong. In Press. Sample size requirements for detecting time-area variations in length composition of tuna purse seine catches. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-SWFC.




Figure 1. South Pacific Treaty region and statistical areas used for sampling of catches and fish sizes.

5557

MONTH

9

[illegible]

Figure 2. Purse Seine Vessel Catch Report (PSVCR) Form used to record daily fishing information. The form is completed by the vessel captains and is submitted to the FFA as required under the Treaty.

ATTC FORM 2 REVISED 8/77

Figure 3. Logbook Abstracting (LA) Form used to record information for selecting wells for sampling of fish sizes.

PURSE SEINE UNLOADING LOGSHEET

VESSEL NAME _____ RADIO CALL SIGN OR
REGIONAL REGISTER NO. _____

(1) PORT _____

OR, IF AT SEA, POSITION: LAT. _____ LONG. _____

(2) DATES

(a) AT UNLOADING POINT

ARRIVAL _____ DEPARTURE _____

(b) AT UNLOADING

COMMENCEMENT _____ COMPLETION _____

(3) PARTIAL OR COMPLETE UNLOADING _____

(4) UNLOADING TO _____

(5) (a) CARRIER VESSEL NAME _____

and RADIO CALL SIGN OR REGIONAL REGISTER NO. _____

or

(b) NAME AND ADDRESS OF COMPANY ACCEPTING FISH _____

(6) DESTINATION OF FISH _____

(7) QUANTITY UNLOADED

	YELLOWFIN	SKIPJACK	BIGEYE	MARLIN	OTHER	UNIT OF MEASUREMENT
ACCEPTED	_____	_____	_____	_____	_____	_____
REJECTED	_____	_____	_____	_____	_____	_____

SIGNATURES

VESSEL MASTER RECEIVING AGENT

Figure 4. Purse Seine Unloading Logsheets (PSUL) used to record cannery unloading weights. The form is completed by the vessel captain and is submitted to the FFA as required under the Treaty.

LENGTH - FREQUENCY SAMPLING STATUS BOARD

MONTH		AREA STRATUM				
		I	II	III	IV	V
January	Catch					
	# Samples					
February	Catch					
	# Samples					
March	Catch					
	# Samples					
April	Catch					
	# Samples					
May	Catch					
	# Samples					
June	Catch					
	# Samples					
July	Catch					
	# Samples					
August	Catch					
	# Samples					
September	Catch					
	# Samples					
October	Catch					
	# Samples					
November	Catch					
	# Samples					
December	Catch					
	# Samples					

Figure 6. Length-Frequency Sampling Status Board used to monitor the size sampling by month-area strata.

VESSEL LENGTH-FREQUENCY SAMPLING PLAN

VESSEL NAME _____ GEAR _____

FORM NO. _____ DATE _____ CANNERY _____

NAME OF FISHING CAPTAIN _____ NAVIGATOR _____

HAILED TONNAGE	
1940	1,000
1941	1,000
1942	1,000
1943	1,000
1944	1,000
1945	1,000
1946	1,000
1947	1,000
1948	1,000
1949	1,000
1950	1,000
1951	1,000
1952	1,000
1953	1,000
1954	1,000
1955	1,000
1956	1,000
1957	1,000
1958	1,000
1959	1,000
1960	1,000
1961	1,000
1962	1,000
1963	1,000
1964	1,000
1965	1,000
1966	1,000
1967	1,000
1968	1,000
1969	1,000
1970	1,000
1971	1,000
1972	1,000
1973	1,000
1974	1,000
1975	1,000
1976	1,000
1977	1,000
1978	1,000
1979	1,000
1980	1,000
1981	1,000
1982	1,000
1983	1,000
1984	1,000
1985	1,000
1986	1,000
1987	1,000
1988	1,000
1989	1,000
1990	1,000
1991	1,000
1992	1,000
1993	1,000
1994	1,000
1995	1,000
1996	1,000
1997	1,000
1998	1,000
1999	1,000
2000	1,000
2001	1,000
2002	1,000
2003	1,000
2004	1,000
2005	1,000
2006	1,000
2007	1,000
2008	1,000
2009	1,000
2010	1,000
2011	1,000
2012	1,000
2013	1,000
2014	1,000
2015	1,000
2016	1,000
2017	1,000
2018	1,000
2019	1,000
2020	1,000
2021	1,000
2022	1,000
2023	1,000
2024	1,000
2025	1,000
2026	1,000
2027	1,000
2028	1,000
2029	1,000
2030	1,000
2031	1,000
2032	1,000
2033	1,000
2034	1,000
2035	1,000
2036	1,000
2037	1,000
2038	1,000
2039	1,000
2040	1,000
2041	1,000
2042	1,000
2043	1,000
2044	1,000
2045	1,000
2046	1,000
2047	1,000
2048	1,000
2049	1,000
2050	1,000
2051	1,000
2052	1,000
2053	1,000
2054	1,000
2055	1,000
2056	1,000
2057	1,000
2058	1,000
2059	1,000
2060	1,000
2061	1,000
2062	1,000
2063	1,000
2064	1,000
2065	1,000
2066	1,000
2067	1,000
2068	1,000
2069	1,000
2070	1,000
2071	1,000
2072	1,000
2073	1,000
2074	1,000
2075	1,000
2076	1,000
2077	1,000
2078	1,000
2079	1,000
2080	1,000
2081	1,000
2082	1,000
2083	1,000
2084	1,000

INFORMATION FOR SAMPLES TO BE TAKEN

[illegible][illegible]

Figure 7. Vessel Length-Frequency Sampling Plan used to prioritize size sampling of individual vessel's catches.

LENGTH FREQUENCY SAMPLING FORM

1. VESSEL: _____ 2. RANDOM / SELECT 3. SPECIES: _____ 4. SAMPLE NO.: _____

5. TRANSSHIPMENT: _____ 6. HOLD / WELL / SET: _____ 7. OCEAN: _____

8. DATE: _____ 9. MEASURED BY: _____ 10. TIME: _____

11. TONNAGE: TOTAL _____ YF _____ SJ _____ OTHERS _____

25	66	107	148
26	67	108	149
27	68	109	150
28	69	110	151
29	70	111	152
30	71	112	153
31	72	113	154
32	73	114	155
33	74	115	156
34	75	116	157
35	76	117	158
36	77	118	159
37	78	119	160
38	79	120	161
39	80	121	162
40	81	122	163
41	82	123	164
42	83	124	165
43	84	125	166
44	85	126	167
45	86	127	168
46	87	128	169
47	88	129	170
48	89	130	171
49	90	131	172
50	91	132	173
51	92	133	174
52	93	134	175
53	94	135	176
54	95	136	177
55	96	137	178
56	97	138	179
57	98	139	180
58	99	140	181
59	100	141	182
60	101	142	183
61	102	143	184
62	103	144	185
63	104	145	186
64	105	146	187
65	106	147	188

12. TOTAL NO.: _____

13. PREDORSAL / FORK LENGTH

14. COMMENTS: _____

Figure 8. Length-Frequency (LF) Form (front) used to record sizes of fish, catches and catch location by vessel well.

SPECIES COMPOSITION

Reporter: _____ Species: _____ Sample #: _____

Date: _____ Reported from: _____

Catcher Vessel: _____ Gear: _____

Flag: _____ Date of catch: _____ Area: _____

Transshipment Vessel: _____

Hatch/Well/Set: _____ Sort Type: _____

Unloading Weight (S/T): _____ Size Code: _____

Total: _____ YF: _____ SJ: _____

Other: _____

Species	Number of Fish	Freq. Sample Number	Biolo. Sample Number	
Yellowfin				
Skipjack				
Bigeye				
Albacore				
Bluefin				
Blk. Skipjack				
Others				
UNK.(YE/BE)				
Total				

Comments: _____

LF0002

Figure 9. Species Composition (SC) Form used to record proportions of each species in the catch by vessel well.

TAG RECOVERY RECORD

Finder _____	
Vessel _____	Gear _____
Area of Recapture _____	Date of Recapture <u> </u> / <u> </u> / <u> </u> Day Mo. Yr.
Tag Number _____	Well _____
<u>color</u> _____	
Species _____	Date Found <u> </u> / <u> </u> / <u> </u> Day Mo. Yr.
Length/Weight _____	City Found <u> </u> / <u> </u> / <u> </u> Day Mo. Yr.
<u>F</u> <u>S</u>	
Place Found _____	
Samples Collected _____	otoliths ☐ Spines ☐
Date Sampled <u> </u> / <u> </u> / <u> </u>	stomach ☐
Fish Number	
Place Mailed _____	Date Mailed <u> </u> / <u> </u> / <u> </u> Day Mo. Yr.
Reward Paid _____	Date Paid <u> </u> / <u> </u> / <u> </u> Day Mo. Yr.
Comments & Notes: _____	Signature _____

Figure 10. Tag Recovery Record used to record weight, length, catch area and other information for tag returns.

For Month of _____

[illegible]

Figure 11. Monthly summary of tags and reward payments form used to account for distribution of rewards.

